



1/15



LINKER LIPID A

FIGURE 1


2/15




LINKER GRAMICIDIN B

FIGURE 2

3/15



MEMBRANE SPANNING LIPID D



MEMBRANE SPANNING LIPID C

FIGURE 3

5/15




FIGURE 5

6/15




FIGURE 6

7/15

08/685329

ADDITION OF 100% (v/v) WHOLE BLOOD TO MONOLAYER MEMBRANES




Figure 7

08/685329

8/15





Figure 8

9/15



Blood
[K⁺] in well

1.34mM

3.35mM

5.03mM

16.75mM

bloodvl.tc
(average 4 electrodes)

Figure 9: Human whole blood K⁺ titration in biosensor membrane.




Figure 10 : 500pM Ferritin response in human whole blood and serum.

11/15

08/685329

ENDOGENOUS SERUM FERRITIN DETECTION FROM 4 DIFFERENT VOLUNTEERS





Figure 11: Ferritin titration curve based on endogenous ferritin levels in human serum.

12/15

08/685329

**METHOD: 5/4 membranes**


1st layers: 10mM GMPE
 1mM DLP
 10uM MSLXXB
 0.8mM MAAD
 0.1uM GaYYSS

2nd layer: 28mM GMPE
 0.28uM GaX..B chosen from the following:
 GaXB
 GaXOB
 GaXOOB
 Ga(XXXB)2

Figure 12

13/15


Figure 13

EXAMPLE OF T_4 ASSAY WITH AMBRI BIOSENSOR

14/15

08/685329


Figure 14



(1) before the addition of gramicidin/SDS

(2) after the addition of gramicidin/SDS; streptavidin and biotinylated Fab'

(3) after the addition of ferritin

Figure 15

(1) before addition of gramicidin/streptavidin

(2) after addition of gramicidin/streptavidin and biotinylated Fab's

(3) after addition of TSH